A TVERBERG-TYPE RESULT ON MULTICOLORED SIMPLICES

JÁNOS PACH

ABSTRACT. Let $P_1, P_2, \ldots, P_{d+1}$ be pairwise disjoint n-element point sets in general position in d-space. It is shown that there exist a point O and suitable subsets $Q_i \subseteq P_i$ $(i = 1, 2, \ldots, d+1)$ such that $|Q_i| \geq c_d|P_i|$, and every d-dimensional simplex with exactly one vertex in each Q_i contains O in its interior. Here c_d is a positive constant depending only on d.

1. INTRODUCTION

Let $P_1, P_2, \ldots, P_{d+1}$ be pairwise disjoint n-element point sets in general position in Euclidean d-space $\Re^d$. If two points belong to the same P_i, then we say that they are of the same *color*. A d-dimensional simplex is called *multicolored*, if it has exactly one vertex in each P_i $(i = 1, 2, \ldots, d+1)$. Answering a question of Bárány, Füredi, and Lovász [BFL90], Vrećica and Živaljević [ZV92], proved the following Tverberg-type result. For every k, there exists an integer $n(k, d)$ such that if $n \geq n(k, d)$, then one can always find k multicolored vertex disjoint simplices with an interior point in common. (For some special cases, see [BL92], [JS91], [VZ94].) This theorem can be used to derive a nontrivial upper bound on the number of different ways one can cut a finite point set into two (roughly) equal halves by a hyperplane.

The aim of this note is to strengthen the above result by showing that there exist "large" subsets of the sets P_i such that *all* multicolored simplices induced by them have an interior point in common.

Theorem. *There exists $c_d > 0$ with the property that for any disjoint n-element point sets $P_1, P_2, \ldots, P_{d+1} \subset \Re^d$ in general position, one can find a point O and suitable subsets $Q_i \subseteq P_i$, $|Q_i| \geq c_d|P_i|$ $(i = 1, 2, \ldots, d+1)$ such that every d-dimensional simplex with exactly one vertex in each Q_i contains O in its interior.*

The proof is based on the $k = d + 1$ special case of the Vrećica-Živaljević theorem (see Theorem 2.1). It uses three auxiliary results, each of them interesting on its own right. The first is Kalai's fractional Helly theorem [K84], which sharpens and generalizes some earlier results of Katchalski and Liu [KL79] (see Theorem 2.2). The second is a variation of Szemerédi's regularity lemma for hypergraphs [S78] (Theorem

Supported by NSF grant CCR-94-24398, PSC-CUNY Research Award 663472 and OTKA-4269. This paper was written while the author was visiting MSRI Berkeley, as part of the Convex Geometry Program. Research at MSRI is supported in part by NSF grant DMS-9022140.

2.3), and the third is a corollary of Radon's theorem [R21], discovered and applied by Goodman and Pollack [GPW96] (Theorem 2.4).

In the next section, we state the above mentioned results and also include a short proof of Theorem 2.3, because in its present form it cannot be found in the literature. Our argument is an adaptation of the approach of Komlós and Sós [KS96]. For some similar results, see [C91],[FR92],[KS95]. The proof of the Theorem is given in Section 3. It shows that the statement is true for a constant $c_d > 0$ whose value is triple-exponentially decreasing in d.

2. Auxiliary results

Theorem 2.1. [ZV92] *Let $A_1, A_2, \dots, A_{d+1}$ be disjoint $4d$-element sets in general position in d-space. Then one can find $d+1$ vertex disjoint simplices with a common interior point such that each of them has exactly one vertex in every A_i, $1 \le i \le d+1$.*

A family of sets is called *intersecting* if they have an element in common.

Theorem 2.2. [K84] *For any $\alpha > 0$, there exists $\beta = \beta(\alpha, d) > 0$ satisfying the following condition. Any family of N convex sets in d-space, which contains at least $\alpha\binom{N}{d+1}$ intersecting $(d+1)$-tuples, has an intersecting subfamily with at least βN members.*

In fact, if N is sufficiently large, then Theorem 2.2 is true for any $\beta < 1 - (1-\alpha)^{1/(d+1)}$. In particular, it holds for $\beta = \alpha/(d+1)$.

Let $\mathcal{H}$ be a $(d+1)$-partite hypergraph whose vertex set is the union of $d+1$ pairwise disjoint n-element sets, $P_1, P_2, \dots, P_{d+1}$, and whose edges are $(d+1)$-tuples containing precisely one element from each P_i. For any subsets $S_i \subseteq P_i$ $(1 \le i \le d+1)$, let $e(S_1, \dots, S_{d+1})$ denote the number of edges of $\mathcal{H}$ induced by $S_1 \cup \ldots \cup S_{d+1}$. In this notation, the total number of edges of $\mathcal{H}$ is equal to $e(P_1, \dots, P_{d+1})$.

It is not hard to see that for any sets S_i and for any integers $t_i \le |S_i|$, $1 \le i \le d+1$,

$$\frac{e(S_1, \dots, S_{d+1})}{|S_1| \dots |S_{d+1}|} = \sum \frac{e(T_1, \dots, T_{d+1})}{|T_1| \dots |T_{d+1}|} \Big/ \binom{|S_1|}{t_1} \cdots \binom{|S_{d+1}|}{t_{d+1}}, \tag{1}$$

where the sum is taken over all t_i-element subsets $T_i \subseteq S_i$, $1 \le i \le d+1$.

Theorem 2.3. *Let $\mathcal{H}$ be a $(d+1)$-partite hypergraph on the vertex set $P_1 \cup \ldots \cup P_{d+1}$, $|P_i| = n$ $(1 \le i \le d+1)$, and assume that $\mathcal{H}$ has at least βn^{d+1} edges for some $\beta > 0$. Let $0 < \varepsilon < 1/2$.*

Then there exist subsets $S_i \subseteq P_i$ of equal size $|S_i| = s \ge \beta^{1/\varepsilon^{2d}} n$ $(1 \le i \le d+1)$ such that
(i) $e(S_1, \dots, S_{d+1}) \ge \beta s^{d+1}$,
(ii) $e(Q_1, \dots, Q_{d+1}) > 0$ for any $Q_i \subseteq S_i$ with $|Q_i| \ge \varepsilon s$ $(1 \le i \le d+1)$.

Proof: Let $S_i \subseteq P_i$ $(1 \le i \le d+1)$ be sets of equal size such that

$$\frac{e(S_1, \dots, S_{d+1})}{|S_1|^{d+1-\varepsilon^{2d}}}$$

is *maximum*, and denote $|S_1| = \ldots = |S_{d+1}|$ by s.

For this choice of S_i, condition (i) in the theorem is obviously satisfied, because

$$\frac{e(S_1, \dots, S_{d+1})}{|S_1|^{d+1-\varepsilon^{2d}}} \ge \frac{e(P_1, \dots, P_{d+1})}{n^{d+1-\varepsilon^{2d}}} = \frac{\beta}{n^{-\varepsilon^{2d}}} \ge \frac{\beta}{s^{-\varepsilon^{2d}}}.$$

Taking into account the trivial relation

$$\frac{e(S_1, \dots, S_{d+1})}{|S_1|^{d+1-\varepsilon^{2d}}} \le s^{\varepsilon^{2d}},$$

the above inequalities also yield that $s \ge \beta^{1/\varepsilon^{2d}} n$.

It remains to verify (ii). To simplify the notation, assume that εs is an integer, and let Q_i be any εs-element subset of S_i $(1 \le i \le d+1)$. Then

$$\begin{aligned} e(Q_1, \dots, Q_{d+1}) = e(&S_1, \dots, S_{d+1}) \\ &- e(S_1 - Q_1, S_2, S_3, ..., S_{d+1}) \\ &- e(Q_1, S_2 - Q_2, S_3, ..., S_{d+1}) \\ &- e(Q_1, Q_2, S_3 - Q_3, ..., S_{d+1}) \\ &\dots \\ &- e(Q_1, Q_2, Q_3, ..., S_{d+1} - Q_{d+1}). \end{aligned}$$

In view of (1), it follows from the maximal choice of S_i that

$$\begin{aligned} &e(S_1 - Q_1, S_2, ..., S_{d+1}) \\ &\quad = (1-\varepsilon)s^{d+1} \frac{e(S_1 - Q_1, S_2, ..., S_{d+1})}{|S_1 - Q_1||S_2|...|S_{d+1}|} \\ &\quad = (1-\varepsilon)s^{d+1} \sum_{\substack{T_i \subseteq S_i, |T_i| = (1-\varepsilon)s \\ 2 \le i \le d+1}} \frac{e(S_1 - Q_1, T_2, ..., T_{d+1})}{[(1-\varepsilon)s]^{d+1}} \Big/ \binom{s}{\varepsilon s}^d \\ &\quad \le (1-\varepsilon)s^{d+1} \frac{e(S_1, S_2, ..., S_{d+1})}{s^{d+1-\varepsilon^{2d}}} [(1-\varepsilon)s]^{-\varepsilon^{2d}} \\ &\quad = e(S_1, ..., S_{d+1})(1-\varepsilon)^{1-\varepsilon^{2d}}. \end{aligned}$$

Similarly, for any $i,\ 2 \le i \le d+1$, we have

$$e(Q_1, ..., Q_{i-1}, S_i - Q_i, S_{i+1}, ..., S_{d+1}) \leq e(S_1, ..., S_{d+1})\varepsilon^{i-1-\varepsilon^{2d}}(1-\varepsilon).$$

Summing up these inequalities, we obtain

$$e(Q_1, \dots, Q_{d+1}) \geq e(S_1, \dots, S_{d+1})(1-(1-\varepsilon)^{1-\varepsilon^{2d}} - \sum_{i=2}^{d+1} \varepsilon^{i-1-\varepsilon^{2d}}(1-\varepsilon))$$
$$\geq e(S_1, \dots, S_{d+1})(1-(1-\varepsilon)^{1-\varepsilon^{2d}} - \varepsilon^{1-\varepsilon^{2d}} + \varepsilon^{d+1-\varepsilon^{2d}}) > 0,$$

as required. □

A $(d+1)$-tuple of convex sets in d-space is called *separated* if any j of them can be strictly separated from the remaining $d+1-j$ by a hyperplane, $1 \leq j \leq d$. An arbitrary family of at least $d+1$ convex sets in d-space is *separated* if every $(d+1)$-tuple of it is separated.

Theorem 2.4. [GPW96] *A family of convex sets in d-space is separated if and only if no $d+1$ of its members can be intersected by a hyperplane.*

Let $n \geq d+1$. Two sequences of points in d-space, $(p_1, \dots, p_n)$ and $(q_1, \dots, q_n)$, are said to have the same *order type* if for any integers $1 \leq i_1 < \dots < i_{d+1} \leq n$, the simplices $p_{i_1} \dots p_{i_{d+1}}$ and $q_{i_1} \dots q_{i_{d+1}}$ have the same orientation [GP93]. It readily follows from the last result that if $C_1, \dots, C_n$ form a separated family of convex sets, then the order type of $(p_1, \dots, p_n)$ will be the same for every choice of elements $p_i \in C_i$, $1 \leq i \leq n$.

3. Proof of Theorem

Let $P_1, \dots, P_{d+1}$ be pairwise disjoint n-element point sets in general position in d-space. If a simplex has precisely one vertex in each P_i, we call it *multicolored*. The number of multicolored simplices is $N = n^{d+1}$.

By Theorem 2.1, any collection of $4d$-element subsets $A_i \subseteq P_i$, $1 \leq i \leq d+1$, induce $d+1$ vertex disjoint multicolored simplices with a common interior point. Thus, the total number of intersecting $(d+1)$-tuples of multicolored simplices is at least

$$\frac{\binom{n}{4d}^{d+1}}{\binom{n-d-1}{3d-1}^{d+1}} > \frac{1}{(5d)^{d^2}}\binom{N}{d+1}.$$

Hence, we can apply Theorem 2.2 with $\alpha = 1/(5d)^{d^2}$. We obtain that there is a point O contained in the interior of at least $\beta N = \beta(1/(5d)^{d^2}, d)n^{d+1}$ multicolored simplices.

Let $\mathcal{H}$ denote the $(d+1)$-partite hypergraph on the vertex set $P_1 \cup \ldots \cup P_{d+1}$, whose edge set consists of all multicolored $(d+1)$-tuples that induce a simplex containing O in its interior.

Set $\varepsilon = 1/2^{d2^d}$, and apply Theorem 2.3 to the hypergraph $\mathcal{H}$ to find $S_i \subseteq P_i$, $1 \leq i \leq d+1$, meeting the requirements. By throwing out some points from each S_i, but retaining a positive proportion of them, we can achieve that the convex hulls of the sets S_i are separated. Indeed, assume e.g. that there is no hyperplane strictly separating $S_1 \cup \ldots \cup S_j$ from $S_{j+1} \cup \ldots \cup S_{d+1}$. By the *ham-sandwich theorem* [B33], one can find a hyperplane h which simultaneously bisects $S_1, \ldots, S_d$ into as equal parts as possible. Assume without loss of generality that at least half of the elements of S_{d+1} are "above" h. Then throw away all elements of $S_1 \cup \ldots \cup S_j$ that are above h and all elements of $S_{j+1} \cup \ldots \cup S_{d+1}$ that are below h. We can repeat this procedure as long as we find a non-separated $(d+1)$-tuple. In each step, we reduce the size of every set by a factor of at most 2.

Notice that in the same manner we can also achieve that e.g. the $(d+1)$-tuple $\{\{O\}, \text{conv}(S_1), \ldots, \text{conv}(S_d)\}$ becomes separated. In this case, h will always pass through the point O, therefore O will never be deleted.

After at most $(d+2)2^d$ steps we end up with $Q_i \subseteq S_i$, $|Q_i| > \varepsilon s$ $(1 \leq i \leq d+1)$ such that $\{\{O\}, \text{conv}(S_1), \ldots, \text{conv}(S_{d+1})\}$ is a separated family. It follows from the remark after Theorem 2.4 that there are only two possibilities: either every multicolored simplex induced by $Q_1 \cup \ldots \cup Q_{d+1}$ contains O in its interior, or none of them does. However, this latter option is ruled out by part (ii) of Theorem 2.3. This completes proof. □

Instead of applying Theorem 2.2, we could have started the proof by referring to the following result of Alon, Bárány, Füredi, and Kleitman [ABFK92], which is also based on Theorem 2.1. For any $\beta > 0$ there is a $\beta_d' > 0$ such that any family of βn^{d+1} simplices induced by n points in d-space has at least $\beta_d' n^{d+1}$ members with non-empty intersection.

Our proof easily yields the following.

Theorem 3.1. *For any $\beta > 0$ there is a $\beta_d'' > 0$ with the property that given any family of βn^{d+1} simplices induced by an n-element set $P \subset \Re^d$, one can find a point O and suitable subsets $Q_i \subseteq P_i$, $|Q_i| \geq \beta_d''|P_i|$ $(i = 1, 2, \ldots, d+1)$ such that every d-dimensional simplex with exactly one vertex in each Q_i contains O.*

Acknowledgement. I am grateful to Imre Bárány, Géza Tóth, and Pavel Valtr for their valuable suggestions.

References

[ABFK92] N. Alon, I. Bárány, Z. Füredi, and D. Kleitman, Point selections and weak ε-nets for convex hulls *Combin. Probab. Comput.* **1**, 189–200.

[BFL90] I. Bárány, Z. Füredi, and L. Lovász, On the number of halving planes, *Combinatorica* **10**, 175–183.

[BL92] I. Bárány and D. Larman, A colored version of Tverberg's theorem, *J. London Math. Soc. (2)* **45**, 314–320.

[B33] K. Borsuk, Drei Sätze über die n-dimensionale euklidische sphäre, *Fundamenta Math.* **20**, 177–190.

[C91] F. R. K. Chung, Regularity lemmas for hypergraphs and quasi-randomness, *Random Structures and Algorithms* **2**, 241–252.

[FR92] P. Frankl and V. Rödl, The uniformity lemma for hypergraphs, *Graphs and Combinatorics* **8**, 309–312.

[GPW93] J. E. Goodman, R. Pollack, and R. Wenger, Geometric transversal theory, in: *New Trends in Discrete and Computational Geometry (J. Pach, ed.), Springer-Verlag, Berlin*, 163–198.

[GPW96] J. E. Goodman, R. Pollack, and R. Wenger, Bounding the number of geometric permutations induced by k-transversals, *J. Combin. Theory, Ser. A,* to appear.

[JS91] J. Jaromczyk and G. Świątek, The optimal constant for the colored version of Tverberg's theorem, *manuscript.*

[K84] G. Kalai, Intersection patterns of convex sets, *Israel J. Math.* **48**, 161–174.

[KL79] M. Katchalski and A. Liu, A problem of geometry in $\Re^d$, *Proc. Amer. Math. Soc.* **75**, 284–288.

[KS95] J. Komlós and M. Simonovits, Szemerédi's regularity lemma and its applications, to appear in: *Paul Erdős is Eighty, Proc. Colloq. Math. Soc. J. Bolyai.*

[KS96] J. Komlós and V. T. Sós, Regular subgraphs of graphs, manuscript.

[R21] J. Radon, Mengen konvexer Körper, die einen gemeinsamen Punkt enthalten, *Math. Ann.* **83**, 113–115.

[S78] E. Szemerédi, Regular partitions of graphs, in: *Problèmes combinatoires et théorie des graphes, Colloq. Internat. CNRS* **260**, CNRS, Paris, 399–401.

[T66] H. Tverberg, A generalization of Radon's theorem, *J. London Math. Soc.* **41**, 123–128.

[VZ94] S. T. Vrećica and R. T. Živaljević, New cases of the colored Tverberg theorem, in: *Jerusalem Combinatorics '93, Contemp. Math.* **178**, Amer. Math. Soc., Providence, 325–334.

[ZV92] R. T. Živaljević and S. T. Vrećica, The colored Tverberg's problem and complexes of injective functions, *J. Combin. Theory Ser. A* **61**, 309–318.

JÁNOS PACH, CITY COLLEGE, CUNY AND, COURANT INSTITUTE, NYU, 251 MERCER STREET, NEW YORK, NY 10012

E-mail address: pachcims6.nyu.edu